

An algorithmic approach to simulate Hamiltonian dynamics

Ashok Ajoy,^{1,2,*} Rama Koteswara Rao,^{2,3} Anil Kumar,^{2,3} and Pranaw Rungta⁴

¹*Birla Institute of Technology and Science - Pilani, Zuarinagar, Goa - 403726, India.*

²*NMR Research Centre, Indian Institute of Science, Bangalore - 560012, India.*

³*Department of Physics, Indian Institute of Science, Bangalore - 560012, India.*

⁴*Indian Institute of Science Education and Research Mohali, Sector-26 Chandigarh, India*

We propose an iterative algorithm to simulate any n -qubit Hamiltonian dynamics. The simulation entails decomposing a unitary operation into a product of simpler unitary operations, where each unitary operation can be easily implemented in a quantum simulator of choice. Our algorithm exploits the completeness of the chosen operator basis to intuitively understand the structure of the system dynamics, and scaling properties of the simulation. We illustrate the algorithm by simulating a unitary that transports quantum information from one end of a n -qubit chain to the other driven by an Ising Hamiltonian (an operation known as quantum state transfer (QST)). This is applied to provide the first experimental verification of the QST protocol by Di Franco et. al. (Phys. Rev. Lett. **101**, 230502 (2008)) in an NMR quantum simulator. We also experimentally test the resilience of this protocol to static errors in the coupling parameters of the simulated Hamiltonian.

PACS numbers: 03.67.Ac, 76.60.-k, 03.67.Lx

Introduction: Feynman [1] alluded to the fact that one could exploit quantum mechanical evolution to simulate the dynamics of another quantum system exponentially faster than in any classical computer. This has been one of the key motivations to build a quantum information processor (QIP) [2]. Simulation not only allows one to study the dynamics of another system, perhaps not naturally available, but also to execute a certain computational task (an algorithm) [3]. More precisely, the simulation problem is the desire to efficiently implement an unitary operation $U = \exp(-i\mathcal{H}t)$ on ones choice of QIP, where $\mathcal{H}$ is an n -qubit Hamiltonian. An efficient simulation is one which requires physical resources that scale polynomially with the problem size n .

The Hamiltonian of any QIP consists of the sum of the intrinsic Hamiltonian $\mathcal{H}_{\text{int}}$, and a part which depends on parameters controlled experimentally by externally applied fields $\mathcal{H}_{\text{ext}}(t)$ [4]. Their specific form, of course, depends upon the inherent physical nature of the QIP. The parameters are chosen such that the following holds to an arbitrary precision:

$$U \approx U_{\text{sim}} = T \exp \left[-i \int_0^\tau dt (\mathcal{H}_{\text{int}} + \mathcal{H}_{\text{ext}}(t)) \right], \quad (1)$$

where τ is a finite time interval. If a QIP allows universal control, i.e in principle it can simulate any desired unitary operation, then there exist the control parameters such that Eq. 1 holds [5]. However, finding them is the real challenge [6]. Instead, U_{sim} is obtained as a sequence (product) of unitaries that are easily implementable in the QIP: $U_{\text{sim}} = U_1 U_2 \dots U_m$ [7]. In general m scales exponentially with the number of qubits n , unless U has additional structure, for instance, its matrix representation is sparse, or it has certain kinds of symmetry with respect to its action on the space of qubits [8]. Hence, the product decomposition of U unravels the physical structure of U : how it entangles different qubits, which part

of the Hilbert space it accesses and how the resources required for its simulation scale with the number of qubits.

Unfortunately, the conventional method to obtain the decomposition does not allow one to probe such structure of U . In the standard methods [9] (for small n) the decomposition is achieved by resolving τ into finer time steps δt_j , with the assumption that $\delta t_j \rightarrow 0$. A first order approximation is invoked, and numerical optimization techniques are employed: A guess decomposition U_g is made, which is then compared with the desired U via an overlap function (like fidelity); U_g is then bought arbitrarily close to U by iteration [9]. Since the process requires infinitesimal time steps δt_j , the decomposition obtained is discontinuous and highly nonintuitive. In this letter, we rectify this problem by algorithmically decomposing U instead into finite time step building blocks that are elements of a complete operator basis [10]. Moreover, the decomposition obtained by our method for small n may allow one to deduce the same for arbitrary n .

In this letter we demonstrate the utility of our algorithm by explicitly determining the decomposition of the unitary that causes *quantum state transfer* (QST) in a linear n -spin chain [11]. QST physically refers to the transport of quantum information from one end of the chain to the other; allowing spin chains to act as analogues of wires in spin-based QIP architectures [12]. QST is slated to play an important role in the initialization of quantum computers, and in linking together the quantum registers of two different spin based QIPs [13]. While QST can be achieved by a hopping transfer between successive spins of the chain, this requires locally addressing each qubit, and is not experimentally viable for large n . Instead, a global control field on the spin chain, which is in general entangling, maybe more efficient [11]. For example, QST can occur if the spins are driven by an Ising Hamiltonian $\mathcal{H}_I$ (Eq. 3 below). But this requires

manufacturing a chain of spins coupled to each other in a very specific way – a task that is fraught with enormous difficulty [13]. So, while several protocols have been suggested for QST using global control Hamiltonians like $\mathcal{H}_{\mathcal{I}}$, there is no experimental means to test their efficiency or their robustness to fabrication errors in the chain. Experimentally simulating the quantum chain allows one to probe such issues.

Here, we apply the decomposition to experimentally simulate QST in an Nuclear Magnetic Resonance (NMR) QIP [14]. We implement the QST protocol recently proposed by Di Franco et al [15] that employs $\mathcal{H}_{\mathcal{I}}$, and does not require the spin chain to be initialized. We find that the resources for simulation scale linearly with the number of qubits, and that the Ising Hamiltonian entangles all the qubits of the chain. The algorithm is also used to obtain decompositions to experimentally *test* the resilience of this QST protocol to errors in the static couplings between spins.

The Quantum State Transfer: Given a n -qubit chain, initially in the state $|\psi_1\rangle \cdots |\psi_n\rangle$, Di Franco et al suggested a protocol [15] by which QST can be achieved by evolving the spin chain under the nearest-neighbour Ising Hamiltonian $\mathcal{H}_I$ (given below) for a time $t_0 = \pi/4J : U_{\text{QST}} = \exp(-it_0\mathcal{H}_I)$. The importance of this protocol is that it bypasses the initialization of the spin medium (except the last qubit); hence the core of the spin chain is not under local control [15].

$$U_{\text{QST}}|\psi_I\rangle|\psi_2 \cdots \psi_{n-1}\rangle|0\rangle \quad (2)$$

$$= \frac{1}{\sqrt{2}} [|0\rangle|\psi_{n-1} \cdots \psi_2\rangle|\psi_I\rangle + i|1\rangle|\psi_{n-1}^\perp \cdots \psi_2^\perp\rangle(I_x|\psi_I\rangle)]$$

where $\langle\psi_i^\perp|\psi_i\rangle = 0 \forall i$. A state locally equivalent to $|\psi_I\rangle$ can be recovered at the end of the chain by measuring the first qubit in the Z basis.

The Ising Hamiltonian $\mathcal{H}_I$ has the form [15],

$$\mathcal{H}_I = \sum_{j=1}^{n-1} J_j I_{jz} I_{(j+1)z} + \sum_{j=1}^n B_j I_{jx}. \quad (3)$$

I 's are a spin-1/2 Pauli matrices and we assume standard commutation relations hold. J_j refers to the inter-neighbour coupling strength, $J_j = 2J\sqrt{4j(n-j)}$; while B_j refers to the coupling strength of a spin site with a local magnetic field, $B_j = 2J\sqrt{(2j-1)(2n-2j+1)}$.

The algorithm: In this letter we illustrate the algorithm by explicitly obtaining a decomposition of the 3-qubit QST unitary: $U_{\text{QST}} = \exp(-it_0\mathcal{H}_I)$, for $t_0 = \pi/4J$. Following the steps outlined here, the algorithm can be easily generalized for any unitary U . We will decompose U_{QST} in the well-known NMR product operator basis [17],

$$\mathbf{B} = \{\mathbf{1}, I_{j\alpha}, 2I_{j\alpha}I_{k\beta}, 4I_{j\alpha}I_{k\alpha}I_{l\gamma}\}, \quad (4)$$

where $\{\alpha, \beta, \gamma\} \in \{x, y, z\}$. The aim of the algorithm is to find the m basis-elements $D_k (\in \mathbf{B})$ and angles θ_k such that,

$$U_{\text{QST}} = \prod_{k=1}^m \exp(-i\theta_k D_k), \quad (5)$$

Since the D_k 's act on different parts of the qubit space, the decomposition reflects the internal structure of U_{QST} in the chosen basis.

Determining D_k 's in Eq. (5) can be looked upon as a search problem, where the search-space is the set $\mathbf{B}$. For an n -qubit system this space is of exponential size 2^{2n} . The central step in the algorithm is to obtain the decomposition of U_{QST} by iteratively reducing this search space (as explained below). By using $\exp(-i\theta_k D_k) = (\cos \frac{\theta_k}{2} \mathbf{1} - i \sin \frac{\theta_k}{2} D_k)$, Eq. (5) can be re-expressed as,

$$U_{\text{QST}} = C_1 \mathbf{1} + C_{D_1} D_1 + \cdots + C_{D_m} D_m + C_{D_1 D_2} D_1 D_2 + \cdots + C_{D_{m-1} D_m} D_{m-1} D_m + \cdots + C_{D_1 \cdots D_m} D_1 \cdots D_m, \quad (6)$$

which can be simplified by using the matrix product operation as the expansion of the unitary in the complete basis $\mathbf{B}$. In the special case that $G_0 = \{\mathbf{1}, D_1, D_2, \cdots, D_m\}$ forms a group, Eq. (6) reduces to,

$$U_{\text{QST}} = C_1 \mathbf{1} + C_{D_1} D_1 + \cdots + C_{D_m} D_m = C_1 \mathbf{1} + \sum_{k=1}^m C_{D_k} D_k, \quad (7)$$

where the coefficients $C_{D_k} = \text{Tr}(U_{\text{QST}} D_k^\dagger) / \text{Tr}(D_k^\dagger D_k)$. Note that G_0 can always be formed by suitable inclusion of elements from $\mathbf{B}$. The closure of the group G_0 implies that in this case *all* the elements D_k in the decomposition of Eq. (5) belong to G_0 . To see this, consider for example the explicit expansion of the U_{QST} as a sum of elements in $\mathbf{B}$:

$$U_{\text{QST}} = \frac{1}{2\sqrt{2}} [1 - iI_{2x} + 2I_{1x}I_{3x} + 2I_{1y}I_{3y} + 2I_{1z}I_{3z} + -i4I_{1x}I_{2x}I_{3x} + i4I_{1y}I_{2x}I_{3y} + i4I_{1z}I_{2x}I_{3z}] \quad (8)$$

The elements that figure in this basis expansion form a group,

$$G_0 = \{\mathbf{1}, I_{2x}, 2I_{1x}I_{3x}, 2I_{1y}I_{3y}, 2I_{1z}I_{3z}, 4I_{1x}I_{2x}I_{3x}, 4I_{1y}I_{2x}I_{3y}, 4I_{1z}I_{2x}I_{3z}\}, \quad (9)$$

Hence, U_{QST} can be decomposed (as in Eq. (5)) by using D_k 's that exclusively belong to G_0 , i.e. the search space for finding D_k is now $G_0 \subset \mathbf{B}$. The closure of the group also provides a means to iteratively reduce this search space: by extinguishing part of U_{QST} to get $U_{\text{QST}}^{(1)}$ so that the only elements in its basis-expansion (similar to Eq. (7)) now form a *subgroup* of G_0 . For instance, consider the unitary operator $U_{\text{QST}}^{(1)}$,

$$U_{\text{QST}}^{(1)} = U_{\text{QST}} \times \exp\left(-i\frac{\pi}{2} 4I_{1y}I_{2x}I_{3y}\right) \quad (10)$$

$$= \frac{1}{2} [1 - iI_{2x} + 2I_{1z}I_{3z} + i4I_{1z}I_{2x}I_{3z}] \quad (11)$$

Now, since $G_1 = \{\mathbf{1}, I_{2x}, 2I_{1z}I_{3z}, 4I_{1z}I_{2x}I_{3z}\}$ is a subgroup of G_0 ; the search-space is reduced to $G_1 \subset G_0 \subset \mathbf{B}$. A similar iterative reduction can be continued by extinguishing part of $U_{\text{QST}}^{(1)}$ to get $U_{\text{QST}}^{(2)}$, so that the search-space G_2 is a subgroup of G_1 , and so on. More generally, let us assume that forming $U_{\text{QST}}^{(1)}$ from U_{QST} requires m' elements,

$$U_{\text{QST}} \times \left(\prod_{k=1}^{m'} \exp(i\theta_k D_k) \right) = U_{\text{QST}}^{(1)}. \quad (12)$$

These m' basis-elements are precisely those that figure in the decomposition of U_{QST} in Eq. (5). We define the norm of the space spanned by the elements of a set G ($G \subset \mathbf{B}$) in a unitary U as,

$$\mathcal{N}_G(U) = \sum_k |C_{D_k}|^2 \text{ where, } D_k \in G \quad (13)$$

With this definition,

$$\mathcal{N}_{G_1}(U_{\text{QST}}) + \mathcal{N}_{(G_0-G_1)}(U_{\text{QST}}) = 1. \quad (14)$$

The role of the m' elements D_k in Eq. (12) is to make $\mathcal{N}_{(G_0-G_1)}(U_{\text{QST}}^{(1)}) = 0$. They way they are ordered (in the decomposition) is chosen by dynamic programming [18]: at each of m' steps one affects a *maximum* reduction in

norm; so that at the end of m' -steps, the norm of the space spanned by $(G_0 - G_1)$ in $U_{\text{QST}}^{(1)}$ is arbitrarily close to 0. For this process, it is sufficient that the D_k 's belong exclusively to $(G_0 - G_1)$. The angles θ_k in Eq. (12) are chosen in such a way that the associated D_k has no support in $U_{\text{QST}}^{(1)}$. Recurrence relations for the dynamic programming process and their scaling will be presented elsewhere [16]. In summary, the 3-qubit U_{QST} has the decomposition:

$$U_{\text{QST}} = \exp\left(i\frac{\pi}{2}4I_{1y}I_{2x}I_{3y}\right) \exp\left(i\frac{\pi}{2}4I_{1z}I_{2x}I_{3z}\right) \exp\left(-i\frac{\pi}{2}I_{2x}\right) \quad (15)$$

The advantage of our algorithmic approach is that by obtaining the decomposition of a unitary for small n , one maybe able to deduce its decomposition for arbitrary n . For U_{QST} , this decomposition is given in Eq. (16), where the upper equation is for odd n , and the bottom one for even n . The number of basis elements in the decomposition (hence the resources required for simulating U_{QST}) scale linearly with n . Since, in general, directed information transport along a quantum chain would involve all the qubits, at least $\mathcal{O}(n)$ resources are needed to simulate it. Our decomposition is hence minimal in this regard. Eq. (16) also reflects the fact that U_{QST} entangles all spins in the chain as suggested by the protocol in Eq. (2).

$$U_{\text{QST}} = \prod_{k=1}^{\lfloor n/2 \rfloor} \exp\left(i\frac{\pi}{2}2^{n-2k+1}I_{ky}I_{(n-k+1)y} \bigotimes_{j=k+1}^{n-k} I_{jx}\right) \exp\left(i\frac{\pi}{2}2^{n-2k+1}I_{kz}I_{(n-k+1)z} \bigotimes_{j=k+1}^{n-k} I_{jx}\right) \times \begin{cases} \exp\left(-i\frac{\pi}{2}I_{(n+1)/2x}\right) \\ \mathbf{1} \end{cases} \quad (16)$$

Experimental Simulation of Ising Hamiltonian: NMR is a pioneering architecture for experimental QIP, where the development of sophisticated control methods over several decades has led to the exploitation of the full range of the system dynamics [6]. In an NMR QIP [17], $\mathcal{H}_{\text{int}} = \frac{1}{2} \sum_{j=1}^n \gamma_j I_{jz} + \frac{1}{2} \sum_{j=2}^n \sum_{k < j} J_{kj} I_{kz} I_{jz}$, where γ_j is the gyromagnetic ratio of the j th qubit (spin), and J_{kj} is the J -coupling between spins j and k . The external control field $\mathcal{H}_{\text{ext}}(t) = u_x(t) \sum_j I_{jx} + u_y(t) \sum_j I_{jy}$ is implemented via radio frequency (RF) pulses (the pulse sequence) that generate single qubit rotations about the x/y -axis—this ensures universal control [5].

In our experiments, we use ^{13}C labeled $^{13}\text{CHFBr}_2$, where ^1H , ^{13}C and ^{19}F are the three-qubits. Experiments were performed at room temperature in a 11.7T magnetic field on an AV-500 spectrometer with a triple resonance QXI probe. The ^1H , ^{13}C and ^{19}F resonance frequencies at this field are 500MHz, 470MHz and 125MHz respectively. The couplings between the three-qubits are

$J_{\text{HC}} = 224.5\text{Hz}$, $J_{\text{HF}} = 49.7\text{Hz}$ and $J_{\text{FC}} = -310.9\text{Hz}$.

Using hard pulses alone or by standard refocusing techniques [17, 19], each element of the product operator basis $\mathbf{B}$ (Eq. (4)) can be implemented experimentally. One can arrive at the pulse sequence that implements U_{QST} by piecing together the pulse sequences to construct all the elements in the decomposition. Moreover, in Eq. (15) (and Eq. (16)) all elements in the decomposition can be implemented by only using nearest neighbour couplings. The other couplings are refocused during the net period of the sequence. Hence an effective qubit chain can be created in the NMR QIP, and the pulse sequence alters the total NMR Hamiltonian so that its evolution is identical to that of the Ising Hamiltonian $\mathcal{H}_I$ after t_0 . We will employ ^1H , ^{13}C and ^{19}F as a 3-qubit chain in that order; the J_{HF} coupling is refocused and not used.

Let $|\psi_I\rangle = \alpha|0\rangle + \beta|1\rangle$ be the state of the first qubit which is to be transferred along the spin chain. The protocol [15] requires that the last qubit be initialized




FIG. 1: (Color online) (a) Pulse sequence for U_{QST} . Filled wide and narrow bars are π and $\pi/2$ pulses respectively. Unfilled bars represent pulses with other flip angles. Dotted lines are erstwhile pulses that have been removed by simplification and can be ignored. The numbers between pulses are delays. The black arrows show the sequences for the individual operators (boxes) in the decomposition of U_{QST} . (b)(i) Theoretical (line) and experimental (points) normalized signal intensity on the third qubit, obtained as the average intensity of the spectral lines corresponding to the state $(|00\rangle + i|11\rangle)|\psi_I\rangle$ (shown in the inset for $\varphi = \pi/2$). Here $f_0 = -40.26\text{kHz}$ (-85.6 ppm). (ii) Experimental signal intensity on the third qubit for different static errors in the Ising Hamiltonian couplings. Two representative pulse sequences used in (ii) are shown in (c), (d). The unfilled bars in (c) and (d) are 76.52° and 168.71° X pulses respectively.

to $|0\rangle$, while the second qubit can be in any arbitrary state. For ease of measurement, we will fix the initial state of the second qubit also to $|0\rangle$. The initial state of the spin chain, $|\psi_I 00\rangle$, was prepared from a $|000\rangle$ pseudopure state using a φ -angle X pulse on the first qubit, where $\varphi = \tan^{-1} \frac{\beta}{\alpha}$. The pulse sequence for U_{QST} (Fig 1(a)) was then applied. However, for the readout of $|\psi_I\rangle$, projective measurement is not possible in an ensemble QIP architecture like NMR. Instead, to verify that QST has indeed occurred, we applied a CNOT(1,3) gate, forcing the system in the state $(|00\rangle + i|11\rangle)|\psi_I\rangle$. The state of the third qubit was measured with a $\bar{Y}$ pulse with the receiver coil in the X direction. The obtained NMR signal shows excellent correlation with $\sin \varphi$ (Fig. 1(b)i), showing that the state $|\psi_I\rangle$ has been transferred to the last qubit.

How does the effectiveness of the QST protocol change if the couplings J_j of the Ising Hamiltonian are imperfect? We consider the case where the deviations of the couplings from their Ising values are equal, i.e $\delta J_1 = \delta J_2$; and use the NMR simulator to test the QST process by determining sequences that simulate the unitaries resulting from the changed Hamiltonians. Two representative examples of these sequences are shown in Fig. 1(c) and 1(d). Not surprisingly, from a decomposition point of view, these sequences have a larger structure compared to the Ising Hamiltonian with perfect couplings. Our experimental results (Fig. 1(b)ii) show that the state recovered is corrupted by an additional phase, and the degree of corruption increases with the error of couplings. Consequently, the final state on the last qubit is not a faithful reflection of $|\psi_I\rangle$, and the QST process is imperfect. Moreover, the transfer efficiency – characterized by the maximum signal amplitude – decreases with the error of couplings.

Summarizing, we have presented an algorithm to provide a decomposition for any desired unitary operator in a manner that provides an intuitive understanding of its internal structure, and which allows the unitary to be simulated by an NMR QIP. We used it to experimentally verify the quantum state transfer protocol [15] in spin chains based on the Ising Hamiltonian.

This letter is dedicated to the memory of Jharana Rani Samal.

* Electronic address: ashok.ajoy@gmail.com

- [1] R. P. Feynman, Int. J. Theor. Phys, **21**, 467 (1982).
- [2] M. A. Nielsen and I. L. Chuang, *Quantum Computation and Quantum Information* (Cambridge University Press, Cambridge, England, 2000).
- [3] E. Farhi and S. Gutmann, Phys. Rev. A, **57**, 2403 (1998).
- [4] N. Khaneja, R. Brockett, S.J. Glaser, Phys. Rev. A, **63** 032308 (2001).
- [5] L. M. K. Vandersypen and I. L. Chuang, Reviews of Mod-

- ern Physics, **76**, 1037 (2004).
- [6] C. A. Ryan, C. Negrevergne, M. Laforest, E. Knill, and R. Laflamme, Phys. Rev. A, **78**, 012328 (2008).
 - [7] J.J. Vartiainen, M. Mottonen, M.M. Salomaa, Phys. Rev. Lett. **92**, 177902 (2004).
 - [8] A.M. Childs, R. Kothari, Quantum Information and Computation, **10**, 669 (2010).
 - [9] N. Khaneja, T. Reiss, C. Kehlet, T. Schulte-Heruggen, S. J. Glaser, J. Magn. Res. **172**, 296 (2005); E.M. Fortunato, M.A. Pravia, N. Boulant, G. Teklemariam, T.F. Havel, D.G. Cory, J. Chem. Phys., **116**, 7599 (2002).
 - [10] A.Y. Kitaev, A. H. Shen, and M. N. Vyalyi, *Classical and Quantum Computation*, Graduate Studies in Mathematics Vol. 47, American Mathematical Society, Providence (2002).
 - [11] M. Christandl, N. Datta, A. Ekert and A.J. Landahl, Phys. Rev. Lett. **92**, 187902 (2004); C. Albanese, M. Christandl, N. Datta and A. Ekert, Phys. Rev. Lett. **93**, 230502 (2004).
 - [12] S.C. Benjamin, S. Bose, Phys. Rev. Lett. **90**, 247901 (2003); S. Bose, Phys. Rev. Lett. **91**, 207901 (2003).
 - [13] A. Kay, arXiv:0903.4274 [quant-ph]; S. Bose, Contemp. Phys. **48**, 13 (2007).
 - [14] J. Zhang, N. Rajendran, D. Suter, Phys. Rev. A. **76**, 012317 (2007); J. Zhang, X. Peng, D. Suter, Phys. Rev. A. **73**, 062325 (2006).
 - [15] C. Di Franco, M. Paternostro and M.S. Kim, Phys. Rev. Lett. **101**, 230502 (2008).
 - [16] A. Ajoy, A. Kumar and P. Rungta (to be published).
 - [17] R.R. Ernst, G. Bodenhausen, and A. Wokaun, *Principles of Nuclear Magnetic Resonance in One and Two Dimensions*, Oxford University Press, Oxford (1987).
 - [18] V.F. Krotov, *Global Methods in Optimal Control*, Marcel Dekker, New York, 1996.
 - [19] H. Yuan, S.J. Glaser, N. Khaneja, Phys. Rev. A **76**, 012316 (2007).